



Contains: 40% [Arrowhead](#), [Blue Sapphire](#), [Concerto](#) and [Nu Blue](#) Kentucky Bluegrass – 60% [Goalkeeper II](#) and [Caddieshack II](#) Perennial Ryegrass

Characteristics

- Adapted to a wide range of turf uses
- Excellent rate of establishment
- Kentucky Bluegrasses have good shade tolerance
- Perennial Ryegrasses have a high level of endophyte enhancement
- Good color and density

Seeding Information

Seeding Rates:

New Seeding: 6 Lbs per 1,000 Sq. Ft.
 Over-seeding: 3 Lbs per 1,000 Sq. Ft.
 Over-seeding High Use Areas: 8 Lbs per 1,000 Sq. Ft. at weekly/monthly intervals as needed

Seeding Dates: March – May, August – October Dormant Seeding: December – March

Germination Times: Perennial Ryegrass – 4-7 Days Kentucky Bluegrass – 10-21 Days

Uses

Elite Sports Mixture is a good quality Kentucky Bluegrass/Perennial Ryegrass mixture that will adapt to a wide range of turf applications. *Elite Sports Mixture* has a good rate of establishment as well as good color and density. An excellent choice for sports fields, but will adapt well on home and commercial lawns, parks, cemeteries and campuses.

This Certified Interagency Blue Tag is your guarantee of variety and superior quality

CERTIFIED SEED

Elite Sports Bluegrass/Ryegrass Mixture
 Interagency Certified Seed

Variety & Kind	Purity	Germin	Origin	Lot Number
Caddieshack II Perennial Ryegrass	29.40%	90%	OR	W8-II-CPRG-032B
Goalkeeper II Perennial Ryegrass	29.40%	90%	OR	846-7-09849
Concerto Kentucky Bluegrass	9.80%	90%	WA	M49-7-302-C
Blue Sapphire Kentucky Bluegrass	9.80%	90%	WA	U21-9-70019-1
Arrowhead Kentucky Bluegrass	9.80%	90%	WA	74.130.06A
Nu Blue Kentucky Bluegrass	9.80%	90%	WA	73.79.06C

Total Pure: 98.00% Crop* 0.05% Inert: 1.94% Weeds: 0.01% AMS: 127
 Net Weight: 50.0 Lbs 22.68 Kgs Tested: JAN-12 Excess Noxious Weeds: None

UNITED SEEDS INC.
 OMAHA 800-456-9901 www.unitedseeds.com
 Des Moines - 800-365-6674

Quality certified by NEBRASKA CROP IMPROVEMENT ASSOCIATION
 MEMBER OF ASSOCIATION OF OFFICIAL SEED CERTIFYING AGENCIES
* The Certifying Agency makes no warranty of any kind, expressly or implied, including merchantability or fitness for purpose, or otherwise, which extends beyond the certification that the seeds inspected met the regulations of this agency. The Seller guarantees this seed to conform to the analysis shown. No further warranty is expressed or implied. Sellers liability is limited to the purchase price of the seed.

Elite Sports

Components and/or percentages of the above blend may vary



General Seeding Guidelines

- A. SEED TIMING
 - a. Seed from April 15 to June 15 or
 - b. August 10 to October 15
 - c. Dormant seeding after November 15 will germinate the following Spring
- B. SITE PREPARATION
 - a. Conduct one soil test per acre to determine soil additives needed
 - b. Ensure soil pH is between 6.0 and 7.0
 - c. Eliminate existing vegetation
 - i. Spray with non-selective herbicide
 - ii. Remove dead vegetation through mechanical means
 - d. Add recommended soil amendments, pH adjustment materials, and fertilizer
 - e. Till soil to a 4-6 inch depth
 - f. Remove rocks and debris larger than one inch diameter
 - g. Pulverize and lightly roll soil
 - h. Apply 1 lb phosphorous (P) (2.27 lbs P₂O₅) / 1000 ft² to soil surface
- C. SEEDING METHOD
 - a. Seeding method is based on slope and soil at site
 - b. Broadcast-, drop-, slit-, or drill-seed flat areas where erosion is not a concern
 - i. Seed should be planted 0.125 to 0.25 inches below soil surface
 - ii. Plant two directions putting ½ of seed down each direction
 - iii. Gently roll or rake seeded area to ensure good soil-to-seed contact
 - c. Hydro-seed steep slopes where erosion is a concern
 - i. When hydro-seeding, broadcast ½ of the seed before hydro-seed mixture is applied and place ½ of the seed in hydro-seed mixture
- D. WATERING REQUIREMENTS
 - a. Water to field capacity immediately after seeding
 - b. The first three weeks after seeding, keep top 1.5 inches of soil moist*
 - c. Weeks four through six after seeding, water 3-4 times per week
 - d. After six weeks, water when grass begins to show draught stress

*Number of times to water per day depends on temperature, humidity, wind, and rainfall
- E. FERTILIZING & MOWING
 - a. Fertilize seeded area with 0.5 lbs nitrogen (N) / 1000 ft² 1, 2, and 3 months after initial seeding, then begin a regular fertilizer schedule
 - b. Mow when grass is one inch longer than desired grass height but do not remove more than 1/3 of grass blade (i.e. mow when grass is 4 inches tall if desired height is 3 inches)