



Contains: 75% [Low Grow Grass Mixture](#) – 25% [Native Wildflower Mixture](#)

Grasses

Whisper Blue Fescue	4.56#
Eco Star Hard Fescue	4.56#
Marco Polo Sheep Fine Fescue	4.56#
Little Bluestem	4.56#
Sideoats Grama	2.88#
Blue Grama	2.88#
Native Wildflowers	8#

Wildflowers

Annual Low Growing Mix	Lead Plant	Purple Prairie Clover
Blackeyed Susan	Maltes Cross	Shasta/Oxeye Daisy
Blanketflower	Mexican Red Hat	Spiked Gayfeather
Blue Flax	Missouri Primrose	White Yarrow
Chicory	Perennial Lupine	Upright Prairie Coneflower
False Sunflower	Greyhead Prairie Coneflower	Evening Primrose
Illinois Bundleflower	Purple Coneflower	
Lanceleaf Coreopsis		

Seeding Information

Seeding Rate: 16-32 Lbs per Acre (1/2 to 1 Lb/1,350 Sq. Ft.)
Seeding Dates: April – June Dormant Seed: December – March
Germination Times: Will vary greatly, 7 – 28 Days as a mix.

Uses

Low Grow Grass/Native Wildflower Mixture is the perfect combination of grasses and wildflowers that provide the beauty of the wildflowers accented by the aesthetics of the low growing prairie type grasses. Weed encroachment is the primary problem with wildflowers. The grasses are added to maintain an adequate environment for the wildflowers to grow and flourish and not be crowded out by unwanted weeds and grasses.

Cultural Practices

Prepare a good firm seedbed and seed with a grassland type drill. It is very important that the seed be placed at the proper depth to encourage quick germination and emergence. Seeding time should be the first two weeks of May or when the soil temperature is warm enough to germinate the warm season grasses.

This analysis tag is your verification of variety and quality

Low Grow Grass/Wildflower Mixture					Antelope Brand Native Wildflower Mixture					
by United Seeds Inc.					UNITEED SEEDS INC.					
Variety and Kind	Purity	Germ.	Origin	Lot Numbers	Perennials		Annuals & Biannuals			
Native Wildflower Mixture	22.94%	82%	US	NWF-11	Blackeyed Susan	Purple Coneflower	Creeping Daisy	Corn Poppy		
Whisper Blue Fescue	14.34%	85%	OR	Z1-8-247	Blanketflower	Purple Prairie Clover	Dwarf Godetia	Catchfly		
Beacon Hard Fine Fescue	14.34%	82%	OR	M6-10-3601A	Blue Flax	Shasta/Oxeye Daisy	Clarkia	Cornflower		
Marco Polo Sheep Fine Fescue	14.34%	84%	OR	L113-M20-8-DF290	Chicory	Spiked Gayfeather	Dwarf Plains Coreopsis	Indianblanket		
Blaze Little Bluestem	8.60%	85%	NE	BLZ-40111	False Sunflower	White Yarrow	Chinese Forget-Me-Not	Lemon Mint		
Butte Sideoats Grama	10.04%	84%	NE	G1543	Illinois Bundleflower	Upright Prairie Coneflower	Rocket Larkspur	Partridge Pea		
Blue Grama	7.17%	85%	NE	L8274	Lanceleaf Coreopsis	Evening Primrose	African Daisy	Plains Coreopsis		
Total Purity: 91.75%	Crop: 0.20%	Inert: 0.02%	Weeds: 0.03%	AMS: 127	Lead Plant		California Poppy	Dwarf Red Coreopsis		
New Weight: 32.0 Lbs (14.5) Kgs	Tested: Jan-12		Noxious Weeds: None		Maltese Cross		Birds Eyes	Baby Blue Eyes		
UNITED SEEDS INC.					Antelope Brand					
Seeding Rate: 16-32 Lbs/Acre					Des Moines - 800-456-9901					
					www.unitedseeds.com					
<small>SELLER MAKES NO WARRANTY, EITHER EXPRESSED OR IMPLIED, EXCEPT THAT THE PRODUCT CONFORMS TO THE DESCRIPTION ON THE LABEL. BUYER AGREES THAT SELLER'S LIABILITY UNDER THIS WARRANTY IS LIMITED TO THE AMOUNT OF THE PURCHASE PRICE AND SELLER HAS NOT RESPONSIBILITY FOR SPECIAL CONSEQUENTIAL OR CONTINGENT DAMAGES.</small>										
					402-331-4800		7500 Burlington, Omaha NE 68127		800-456-9901	

Components and/or percentages of the above blend may vary



General Seeding Guidelines

A. SEED TIMING

- a. Seed from April 15th to June 21st or
- b. Dormant Seeding after November 15th will germinate the following Spring and requires at least 25% more seed because some will be lost

B. SITE PREPARATION

- a. Conduct one soil test per acre to determine soil additives needed
- b. Ensure soil pH is between 6.0 and 7.0
- c. Eliminate existing vegetation
 - i. Spray with non-selective herbicide
 - ii. Remove dead vegetation through mechanical means
- d. Add recommended soil amendments, pH adjustment materials, and fertilizer
- e. Plow and disk area twice or till soil to a 4-6 inch depth
- f. Remove rocks and debris larger than one inch diameter
- g. Culti-pack to ensure a firm seedbed
- h. Apply 1 lb phosphorous (P) (2.27 lbs P₂O₅) / 1000 ft² to soil surface

C. SEEDING METHOD

- a. Seeding method is based on slope and soil at site
- b. Broadcast-, drop-, slit-, or drill-seed flat areas where erosion is not a concern
 - i. Seed should be planted 0.125 to 0.25 inches below soil surface
 - ii. Plant two directions putting ½ of seed down each direction
 - iii. Gently roll, rake, or culti-pack seeded area to ensure good soil-to-seed contact
- c. Hydro-seed steep slopes where erosion is a concern
 - i. When hydro-seeding, broadcast ½ of the seed before hydro-seed mixture is applied and place ½ of the seed in hydro-seed mixture

D. WATERING REQUIREMENTS

- a. Water to field capacity immediately after seeding
 - b. The first three weeks after seeding. keep top 1.5 inches of soil moist*
 - c. Weeks four through six after seeding, water 3-4 times per week
 - d. Six weeks after seeding water when plants begin to show draught stress
- *Number of times to water per day depends on temperature, humidity, wind, and rainfall

E. FERTILIZING & MOWING

- a. Fertilization schedule should be based on soil tests for native grasses
- b. Mowing is important for native grass establishment
 - i. Four weeks after seeding mow to a height of 6 inches
 - ii. Following first mowing, mow every 4th weeks at a height of 12 inches until August and then cease mowing
 - iii. Do not mow after the first year